

## High-pressure spectroscopic study of siderite (FeCO<sub>3</sub>) with a focus on spin crossover

VALERIO CERANTOLA<sup>1,\*</sup>, CATHERINE MCCAMMON<sup>1</sup>, ILYA KUPENKO<sup>1,2</sup>, INNOKENTY KANTOR<sup>2</sup>,  
CARLO MARINI<sup>2</sup>, MAX WILKE<sup>3</sup>, LEYLA ISMAILOVA<sup>1</sup>, NATALIA SOLOPOVA<sup>1</sup>,  
ALEKSANDR CHUMAKOV<sup>2</sup>, SAKURA PASCARELLI<sup>2</sup> AND LEONID DUBROVINSKY<sup>1</sup>

<sup>1</sup>Bayerisches Geoinstitut, Universität Bayreuth, D-95440 Bayreuth, Germany

<sup>2</sup>European Synchrotron Radiation Facility, 38043 Grenoble, France

<sup>3</sup>Deutsch GeoForschungsZentrum GFZ, D-14473 Potsdam, Germany

### ABSTRACT

Fe-bearing carbonates have been proposed as possible candidate host minerals for carbon inside the Earth's interior and hence their spectroscopic properties can provide constraints on the deep carbon cycle. Here we investigate high-pressure spin crossover in synthetic FeCO<sub>3</sub> (siderite) using a combination of Mössbauer, Raman, and X-ray absorption near edge structure spectroscopy in diamond-anvil cells. These techniques sensitive to the short-range atomic environment show that at room temperature and under quasi-hydrostatic conditions, spin crossover in siderite takes place over a broad pressure range, between 40 and 47 GPa, in contrast to previous X-ray diffraction data that described the transition as a sharp volume collapse at approximately 43 GPa. Based on these observations we consider electron spin pairing in siderite to be a dynamic process, where Fe atoms can be either high spin or low spin in the crossover region. Mode Grüneisen parameters extracted from Raman spectra collected at pressures below and above spin crossover show a drastic change in stiffness of the Fe-O octahedra after the transition, where they become more compact and hence less compressible. Mössbauer experiments performed on siderite single crystals as well as powder samples demonstrate the effect of differential stress on the local structure of siderite Fe atoms in a diamond-anvil cell. Differences in quadrupole splitting values between powder and single crystals show that local distortions of the Fe site in powder samples cause spin crossover to start at higher pressure and broaden the spin crossover pressure range.

**Keywords:** Siderite, carbonate, deep carbon cycle, spin transition, diamond-anvil cell, Mössbauer spectroscopy, Raman spectroscopy, XANES, differential stress

### INTRODUCTION

Chemical and physical interactions between atmosphere, biosphere, and geosphere are well known and established processes on our planet. However, the mechanisms governing the dynamics and the stability of materials in the deep Earth's interior are still the subject of ongoing debate. The possibility of carbon cycling through the deep Earth (e.g., Dasgupta and Hirschmann 2010) is demonstrated by observations such as the occurrence of diamonds from the lower mantle, carbonate inclusions in mantle xenoliths and diamonds, and the presence of CO<sub>2</sub> in gases from volcanic eruptions (e.g., Brenker et al. 2007; Harte 2011; Walter et al. 2011). However, the size of the carbon reservoir inside the Earth is still not well constrained, and is closely linked to the nature of the dominant carrier(s) of carbon down to the core-mantle boundary (e.g., Biellmann et al. 1993; Lavina et al. 2009; Stagno et al. 2011; Boulard et al. 2012).

Deep carbon is predominantly stored in accessory phases as a consequence of its low solubility in dominant mantle minerals (e.g., Keppler et al. 2003), where these accessory phases include carbonates, diamonds/graphite, methane, and carbides, depending on pressure, temperature, and oxygen fugacity. In highly reducing environments (i.e., low oxygen fugacity), the crystal-

line form of carbon is graphite or diamond. At more oxidizing conditions, carbonates are favored due to the reaction between elemental carbon and oxygen to form (CO<sub>3</sub>)<sup>2-</sup> groups that bond to other cations such as Ca<sup>2+</sup>, Mg<sup>2+</sup>, Fe<sup>2+</sup>, Ni<sup>2+</sup>, and Na<sup>+</sup> depending on the composition of the original bulk assemblage.

The three major carbonate components in the crust and upper mantle are CaCO<sub>3</sub> (calcite), MgCO<sub>3</sub> (magnesite), and FeCO<sub>3</sub> (siderite). The presence of carbonates in the Earth's interior is related to the subduction process, one of the first steps in cycling carbon through the Earth. The CaCO<sub>3</sub>-MgCO<sub>3</sub>-FeCO<sub>3</sub> system has been experimentally investigated at relatively low pressures and moderate temperatures ( $P < 3.5$  GPa,  $T < 1100$  °C) since many decades (e.g., Goldsmith et al. 1962; Rosenberg 1967; Merrill and Bassett 1975; Franzolin et al. 2011). Low-pressure ternary phase diagrams indicate the formation of extensive solid solutions, with compositions depending on pressure and temperature. Observations from geophysics (e.g., Fukao et al. 2009) and natural samples (e.g., Brenker et al. 2007; Walter et al. 2011) suggest that some subducted slabs penetrate the lower mantle. In this case, carbonates could be stable phases due to the proposed higher oxygen fugacity in the region of the slab. The properties of carbonates are, however, not well understood, and many questions remain open regarding their stability, composition, and geophysical properties.

Previous high-pressure studies on the carbonate end-members

\* E-mail: valerio.cerantola@gmail.com

$\text{CaCO}_3$ ,  $\text{MgCO}_3$ , and  $\text{FeCO}_3$  revealed high-pressure phase transitions in all phases (e.g., Ono et al. 2005; Merlini et al. 2012; Biellmann et al. 1993; Isshiki et al. 2004). Magnesite shows a structural change at 115 GPa and 2100–2200 K (Isshiki et al. 2004), whereas magnesiosiderite experiences a volume collapse of almost 10% between 40 and 50 GPa at ambient temperature, caused by a high-spin to low-spin (HS-LS) transition of iron (Lavina et al. 2009). Shi et al. (2008) predicted the spin transition of Fe in pure siderite by ab initio simulations to be between 15 to 28 GPa, while the first experimental observation was reported by Mattila et al. (2007) to occur at roughly 50 GPa from natural siderite powder using X-ray emission spectroscopy. Lavina et al. (2009) described a sharp volume collapse at 43 GPa from X-ray single-crystal diffraction, whereas Farfan et al. (2012) observed spin crossover in a (Mn,Mg)-siderite at 46 GPa using Raman spectroscopy and X-ray single-crystal diffraction. Lin et al. (2012) studied  $\text{Mg}_{0.35}\text{Fe}_{0.65}\text{CO}_3$  up to 80 GPa at ambient temperature and observed the spin transition at 45 GPa, where their data suggested that the low-spin state has different vibrational and elastic properties compared to the high-spin state. Temperature extends the spin crossover region of magnesiosiderite (Liu et al. 2014). Merlini et al. (2012) postulated that the low-spin state could contribute to the stabilization of Fe-bearing Ca,Mg-carbonate (so-called dolomite-III) above 35 GPa and 2000 K, and Liu et al. (2015) reported an orthorhombic phase of magnesiosiderite above 50 GPa and 1400 K that also appeared to be stabilized by spin crossover. It is therefore clear that iron can play a fundamental role in the behavior of carbonates at high pressure and high temperature.

To elucidate the effect of iron on the behavior of carbonates at high pressure, in particular the nature of spin crossover, we undertook a combined spectroscopic study of the siderite end-member. Nuclear resonance techniques (including Mössbauer spectroscopy) provide a sensitive method to detect the response of Fe atoms to physical changes like spin crossover, and X-ray absorption near edge structure (XANES), and Raman spectroscopies provide complementary information.

## METHODOLOGY

### Synthesis

Iron oxalate ( $\text{FeC}_2\text{O}_4$ ) was used as a starting material to synthesize  $\text{FeCO}_3$ . Commercial  $\text{FeC}_2\text{O}_4$  from Alfa Aesar was used for  $^{57}\text{Fe}$ -unenriched samples, while for  $^{57}\text{Fe}$ -enriched samples, iron oxalate was synthesized by chemical reaction of

$^{57}\text{Fe}$  metal with acetic acid, forming  $^{57}\text{Fe}$ -diacetate precipitate,  $^{57}\text{Fe}(\text{C}_2\text{H}_3\text{O}_2)_2$ . The precipitate was then reacted in the presence of oxalic acid to produce  $^{57}\text{Fe}$ -enriched iron oxalate. All chemical reactions were run under argon to prevent oxidation of iron to  $\text{Fe}^{3+}$  at ambient conditions.  $\text{FeC}_2\text{O}_4$  powder was then sealed in gold capsules of 2.3 mm outer diameter and loaded into an externally heated cold seal vessel and run at 2 kbar and 360 °C for 7 days, following French (1971). The obtained powder was used to grow  $\text{FeCO}_3$  single crystals using a multi-anvil press at Bayerisches Geoinstitut. The powder was pressed inside Re capsules to avoid loss of Fe and placed inside assemblies with  $\text{LaCrO}_3$  heaters where temperature was measured using  $\text{W}_{75}\text{Re}_{25}/\text{W}_{97}\text{Re}_3$  thermocouples. The samples were annealed for 10 min at 18 GPa and 1600 °C, which yielded single crystals of approximately 5 to 50  $\mu\text{m}$  diameter. The same procedure was used to synthesize both  $^{57}\text{Fe}$ -enriched (96%  $^{57}\text{Fe}$ ) and unenriched (2%  $^{57}\text{Fe}$ )  $\text{FeCO}_3$  single crystals.

### High-pressure experiments


Diamond-anvil cells (DACs), type BX-90 (Kantor et al. 2012), were mounted with diamonds with culet sizes of 250  $\mu\text{m}$  and rhenium gaskets with 120  $\mu\text{m}$  starting diameter hole were employed in all experiments. We used neon gas loaded under pressure (1.4 kbar) (Kurnosov et al. 2008) as a pressure-transmitting medium to obtain hydrostatic conditions, and we measured the wavelength of the ruby fluorescence before and after each experimental run to calculate pressure (Mao et al. 1986). The difference in values contributed to the pressure uncertainty, which was estimated to be 1 and 3 GPa for single crystal and powder experiments, respectively. All DACs were prepared by loading a ~15–20  $\mu\text{m}$  diameter isomeric  $\text{FeCO}_3$  single-crystal chip and a smaller ruby crystal (Fig. 1a), with the exception of Mössbauer measurements using a radioactive source where more than 10  $^{57}\text{Fe}$ -enriched crystals were used in the pressure chamber to optimize signal acquisition time and signal-to-noise ratio (Fig. 1b). A similar series of Mössbauer measurements was also performed using  $\text{FeCO}_3$  powder, which was compressed to a pellet before loading in the DAC to reduce porosity between the grains. All single crystals survived intact throughout each experimental run without breaking or cracking. To test the degree of hydrostaticity, in some experiments we placed two or three ruby crystals in different positions inside the pressure chamber. In all cases we measured the pressure from each ruby within the same experiment to be within  $\pm 1$  GPa of each other.

## IN SITU ANALYTICAL TECHNIQUES

### Mössbauer spectroscopy

$^{57}\text{Fe}$  Mössbauer spectra were recorded at room-temperature in transmission mode on a constant acceleration Mössbauer spectrometer with a nominal 370 MBq  $^{57}\text{Co}$  high specific activity source in a 12  $\mu\text{m}$  thick Rh matrix (referred to below as “in house” spectra). Spectra collection took 2 to 4 days for each pressure point. A set of Mössbauer spectra was also collected during compression using the Synchrotron Mössbauer Source (SMS) at the European Synchrotron Radiation Facility (ESRF), Grenoble, France, on the Nuclear Resonance beamline (Rüffer

**FIGURE 1.** Photographs of  $\text{FeCO}_3$  samples in DACs at the indicated pressures with (a) one single crystal and (b) multiple single crystals, each surrounded by Ne. Both loadings contain a small ruby crystal as a pressure calibrant. (Color online.)



and Chumakov 1996) ID18. The narrow ( $\sim 6$  neV) energy component of X-rays at the Mössbauer energy of 14.4 keV was extracted from a wide spectrum of synchrotron radiation using a  $^{57}\text{FeBO}_3$  single-crystal monochromator and focused to a beam width of size of  $10 \times 15 \mu\text{m}^2$  using Kirkpatrick-Baez mirrors (Potapkin et al. 2012). Each SMS spectrum took approximately 30 min to collect. The velocity scales of all Mössbauer spectra were calibrated relative to 25  $\mu\text{m}$  thick  $\alpha$ -Fe foil, and all spectra were fitted using the software package MossA (Prescher et al. 2012). All Mössbauer measurements were performed on  $^{57}\text{Fe}$ -enriched  $\text{FeCO}_3$ .

### Raman spectroscopy

Raman measurements were performed on a DILOR XY triple spectrometer using the 514 nm line of  $\text{Ar}^+$  laser with  $2 \text{ cm}^{-1}$  spectral resolution and 2  $\mu\text{m}$  spatial resolution at 300 mW of power. Data acquisition was performed with a frequency range extending from 150 to 1300  $\text{cm}^{-1}$ . To maximize the signal-to-noise ratio, each spectrum was collected in three steps over three smaller regions of frequency, where each region was measured five times using an exposure time of 60 s, and then merged together at the end of the acquisition. Orientation effects of the single crystals and hence any polarization effects on the spectra are considered to be negligible since we used an unpolarized green Raman source,

and all crystals were maintained in the same orientation during the experiment. All Raman data presented in the paper were collected on the same single crystal up to the target pressure. Raman measurements were performed at Bayerisches Geoinstitut, Bayreuth, Germany, and spectral fitting was carried out using the software package PeakFit (Systat Software).

### XANES spectroscopy

Fe  $K$ -edge XANES measurements were performed at ESRF at the energy-dispersive X-ray absorption spectroscopy (XAS) beamline ID24 (e.g., Pascarelli et al. 2006). The beam was focused horizontally using a curved polychromator Si 111 crystal in Bragg geometry and vertically with a bent Si mirror. The obtained cross section is about  $3 \times 5 (\text{H} \times \text{V}) \mu\text{m}^2$ . The measured XANES spectra were normalized using the Athena software package (Ravel and Newville 2005), and the second-order polynomial for the pixel to energy conversion parameters was calibrated using a reference  $\alpha$ -Fe foil spectrum.

## RESULTS

### Mössbauer spectroscopy

Mössbauer spectroscopy provides information about the electronic, magnetic, and structural properties of specific elements within a material. The extremely narrow line width of  $\gamma$  rays (4.5




**FIGURE 2.** Room-temperature Mössbauer spectra of  $\text{FeCO}_3$  (a) single crystals and (c) powder collected in house at the indicated pressures. (b) SMS spectra of one  $\text{FeCO}_3$  single crystal collected at ESRF, Grenoble (France). Solid circles = experimental data; black lines = full transmission integral fit; blue and red lines = individual subspectra. Note that the subspectra do not add up to the total spectrum due to the properties of the full transmission integral fit. Percentage bars indicate the relative absorption. (Color online.)

$\times 10^{-9}$  eV in the case of  $^{57}\text{Fe}$ ) provides Mössbauer spectroscopy with the sensitivity to detect spin transitions in compounds with Mössbauer-active nuclei.

The evolution of  $\text{FeCO}_3$  Mössbauer spectra at 298 K and increasing pressure is similar to spectra already reported for other non-magnetic compounds containing  $\text{Fe}^{2+}$  in an octahedral environment, for example ferropericlaire ( $\text{Fe,MgO}$ ) (Kantor et al. 2006b). At pressures below 40 GPa,  $\text{Fe}^{2+}$  in octahedral coordination is represented by a doublet, while at higher pressures the doublet is progressively replaced by a singlet, which at pressures higher than 45 GPa is the only component present in the spectra of single-crystal  $\text{FeCO}_3$  (Figs. 2a and 2b). Mössbauer spectra of powder  $\text{FeCO}_3$  show a similar evolution with increasing pressure, but over a larger pressure interval (Fig. 2c).

The center shift (CS) is sensitive to the s-electron density at the nucleus, while the quadrupole splitting (QS) measures the electric field gradient (EFG) caused by a non-symmetrical charge distribution around the nucleus. The evolution of hyperfine parameters with increasing pressure for both single crystals and powder samples is shown for CS and QS in Figures 3a and 3b, respectively. The general trend of the CS is to decrease with increasing pressure, where the CS of the singlet is approximately 0.3 mm/s lower than the CS for the doublet (Fig. 3a). The behavior of the CS is similar for single crystals and powder; however for the QS the behavior differs, which will be discussed in more detail below. Here we just mention that the QS of the doublet




**FIGURE 3.** Room-temperature center shift (a) and quadrupole splitting (b) (relative to  $\alpha\text{-Fe}$ ) of  $\text{FeCO}_3$  as a function of pressure. Solid circles = in-house single-crystal experiments; open circles = in-house powder experiment; solid triangles = SMS single-crystal experiment.

for single crystals remains essentially constant with increasing pressure with a slight decrease just before the singlet appears at 45 GPa.

Mössbauer spectra of  $\text{FeCO}_3$  single crystals collected in house and using the SMS are slightly different. Each SMS spectrum shows significantly higher resonant absorption and unequal areas of doublet components compared to spectra collected using a radioactive source (Fig. 4). Both of these features are due to properties of the synchrotron X-ray source, namely the low amount of non-resonant radiation and the polarization of the source, which leads to selective excitations of nuclear levels. However all hyperfine parameters of single-crystal spectra are the same within experimental uncertainty (Figs. 2, 3, and 5), hence SMS and in house spectra can be used interchangeably for interpreting hyperfine parameter variations. In contrast there are significant differences between in house spectra of single crystal and powder  $\text{FeCO}_3$ , including the line widths (Fig. 5), which will be discussed below in the context of the effect of stress in DACs.

### Raman spectroscopy

Raman spectroscopy provides direct insight into the vibrational behavior of ions or groups of ions bonded together,




**FIGURE 4.** Comparison of room-temperature  $\text{FeCO}_3$  Mössbauer spectra: (a) SMS collected on one single crystal at 19 GPa; (b) in house collected on multiple single crystals at 18 GPa; and (c) in house collected on powder at 20.5 GPa. The area asymmetry of the doublet in a is due to polarization of the synchrotron Mössbauer source. Percentage bars indicate the relative absorption. (Color online.)

such as  $(\text{CO}_3)^{2-}$ , and their variation as a response to changes of external parameters such as pressure and temperature. In  $\text{FeCO}_3$ ,  $(\text{CO}_3)^{2-}$  groups form planes perpendicular to the  $c$  axis with Fe occupying the interstitial space between them to form octahedral sites coordinated by oxygen atoms. This structural arrangement is typical for carbonates and is known as the calcite or magnesite-type structure. For symmetry reasons (e.g., Rutt and Nicola 1974) the cations give rise to modes that are Raman inactive. Hence, typical Raman spectra of calcite-type carbonates show mainly external and internal vibrations due to the motion of  $(\text{CO}_3)^{2-}$  ions.

At ambient conditions carbonate spectra show six Raman bands in the region from 200 to  $1800\text{ cm}^{-1}$ . However, above  $1300\text{ cm}^{-1}$  only two bands are detected and correspond to weak asymmetric stretching vibrations of  $(\text{CO}_3)^{2-}$ . We neglect these bands to avoid the signal from the diamonds and focus on frequencies below  $1300\text{ cm}^{-1}$ . In this region, the  $\text{FeCO}_3$  Raman spectrum has four vibrational modes ( $176$ ,  $274$ ,  $728$ , and  $1077\text{ cm}^{-1}$ ) (Fig. 6). Based on previous studies (Popkov et al. 1973; Rutt and Nicola 1974) we assign the two lower frequency lines to external  $E_g$  modes that are translations between cation and anion groups and the two higher frequency lines to internal modes,  $E_g$  and  $A_{1g}$ , which are caused by symmetric bending and stretching of  $(\text{CO}_3)^{2-}$ , respectively. A broad low-intensity peak was also observed in the region around  $500\text{ cm}^{-1}$ , which is attributed to electronic Raman scattering from the  $\text{Fe}^{2+}$  ion (Rutt and Nicola 1974). However, the peak is only visible at ambient conditions and is too weak to be observed at higher pressures.

All bands shift to higher frequencies with increasing pressure up to  $40\text{ GPa}$ , but at slightly higher pressure there is a clear change in the behavior of Raman modes (Figs. 6 and 7). The  $A_{1g}$  mode splits with the growth of a second peak at slightly lower frequencies. With increasing pressure the new Raman band becomes progressively more intense, growing at the expense of the higher frequency peak until the latter completely disappears between  $47$  and  $50\text{ GPa}$ . The internal  $E_g$  line jumps discontinuously to higher wavenumbers (from  $782$  to  $840\text{ cm}^{-1}$ ) at roughly




**FIGURE 5.** Effect of pressure on the full-width at half-maximum (FWHM) of  $\text{FeCO}_3$  spectra. Open circles = in house powder; solid circles = in-house single crystals; solid triangles = SMS single crystal. Dashed lines indicate the pressure at which spin crossover starts for single crystal ( $41\text{ GPa}$ ) and powder ( $52\text{ GPa}$ ) measurements.

$43\text{ GPa}$ , and a broad hump emerges at  $755\text{ cm}^{-1}$ , which might also be attributed to the electronic Raman scattering of the  $\text{Fe}^{2+}$  ions at high pressures. The external  $E_g$  vibrations are also affected by the transition. The  $E_g$  vibrational mode at  $289\text{ cm}^{-1}$  completely disappears after  $45\text{ GPa}$ , while the higher frequency external  $E_g$




**FIGURE 6.** Raman spectra of  $\text{FeCO}_3$  collected at ambient temperature and the indicated pressures. The two modes below  $500\text{ cm}^{-1}$  are external or lattice vibrations of  $\text{CO}_3^{2-}$  relative to Fe atoms, while modes above  $700\text{ cm}^{-1}$  are internal vibrations of  $\text{CO}_3^{2-}$ . The insets indicate the lower frequency Raman bands for the spectra on a different scale.




**FIGURE 7.** Pressure variation of the peak shift of observed Raman modes of  $\text{FeCO}_3$ . Dashed gray area indicates the spin crossover pressure range. The frequency uncertainty is smaller than the size of the symbols.




**FIGURE 8.** Pressure variation of Raman mode intensity normalized to the most intense mode,  $A_{1g}$ , in  $\text{FeCO}_3$ . Solid circles = lower frequency external  $E_g$  vibration; open circles = higher frequency external  $E_g$  vibration; solid triangles = internal  $E_g$  vibration (see also Figs. 7 and 8). Note that the lowest frequency external  $E_g$  mode disappears completely above 45 GPa.

vibration shifts from 449 to 543  $\text{cm}^{-1}$  between 43 and 45 GPa (Figs. 6 and 7). Above 47 GPa and up to 58 GPa, the highest pressure reached by Raman spectroscopy, the new HP Raman modes continues to shift to higher frequencies, although with lower slopes than observed at pressures below the transition.

Normalized spectral intensities highlight the progressive decrease of intensity of lower frequency bands between 0 and 40 GPa (Fig. 8). At higher pressures (>45 GPa) the high frequency external and internal  $E_g$  modes, which are weak and difficult to distinguish in spectra near the transition pressure, gradually become more intense.

### XANES spectroscopy

X-ray absorption fine structure (XAFS) spectroscopy is an element-specific method to study the local atomic structure within a material. XAFS spectra can be divided into different regions based on the energy range from the absorption edge in the spectrum. Our study focuses on the XANES region, which is generally taken to extend 50 eV beyond the edge itself.

Figure 9 shows normalized XANES spectra of  $\text{FeCO}_3$  collected during compression. The 1 bar spectrum is consistent with the data of Wilke et al. (2001), although the pre-edge peak is less pronounced in our study due to differences in energy resolution. Each spectrum is characterized by two peaks (identified in Fig. 9 as “1” and “2”) below 7150 eV. Their relative intensities change at increasing pressure and their positions shift to higher energies (Fig. 10). Peak shifts of the main-edge energy on compression was already observed in previous studies on ferropicroclase (Kantor et al. 2006a; Narygina et al. 2009) and bridgmanite (Narygina et al. 2009) and is related to changes in the electronic structure due to shortening of Fe-O distances. This shortening is directly reflected in the shift of the first EXAFS maximum between 7160 and 7180 eV to higher energy, which may be described by the relation  $\Delta E \times R^2 = \text{const.}$  (e.g., Bianconi et al. 1983; Wilke et al. 2007), where  $\Delta E$  is the energy difference between the onset of the




**FIGURE 9.** (a) Normalized XAS spectra of  $\text{FeCO}_3$  at room temperature collected on compression at the indicated pressures. The inset at the bottom right highlights the pre-edge region of the 1 bar spectrum. (b) Close-up on the evolution of the pre-edge region with pressure. The spectra are shifted vertically for clarity.




**FIGURE 10.** Pressure variation of maximum peak positions for  $\text{FeCO}_3$  XANES spectra. The peaks are labeled according to Figure 9 and the gray dashed area indicates where the transition is observed in the XANES spectra.

edge and the first EXAFS maximum and  $R$  is the Fe-O distance.

All peaks change strongly between 40 and 44 GPa, where the pre-edge region shows the largest variation (Figs. 9a and 10). The pre-edge peak at  $\sim 7112$  eV, which is not well resolved in this data set, becomes more intense and sharper above 44 GPa (Fig. 9b). At  $\sim 7117$  eV,  $\sim 5$  eV higher than the pre-edge, a shoulder present at lower pressure evolves into a better-resolved feature,




**FIGURE 11.**  $\delta\mu/\delta E$  derivatives of room-temperature XANES spectra of  $\text{FeCO}_3$  at the indicated pressures. The vertical unit is arbitrary, and the derivatives are taken from the normalized spectra (Fig. 10). Gray arrows and dashed lines indicate the energies at which changes are observed at 44 GPa.

so that it appears like an additional pre-edge. The gray dashed area in Figure 10 marks the energy where the appearance of this feature is observed, matching the discontinuity in the trends of the main- and pre-edge peaks. Similarly, the main-edge region (7120–7140 eV) shows considerable changes between 40 and 44 GPa. The intensities of the main peaks are reversed for spectra above 40 GPa, where the second peak becomes more intense than the first (Figs. 9–12). Additionally, all spectra above 44 GPa show the presence of a new pronounced feature, identified as “3” in Figure 9.

The derivative  $\delta\mu/\delta E$  provides a more sensitive probe of changes in the spectra (Fig. 11). The gray arrows together with the dashed lines highlight three features that are related to the changes observed in the spectra: (1) the derivative maximum at  $\sim 7115$  eV that disappears below 44 GPa and is substituted by another maximum at  $\sim 7117$  eV; (2) a shoulder at  $\sim 7128$  eV that appears above 44 GPa and sharpens on further increase of pressure to 51.5 GPa; and (3) the derivative minimum at  $\sim 7150$  eV that appears above 44 GPa. These changes in the pre-edge XANES and low-energy EXAFS regions imply modification of the local structure around Fe atoms and/or chemical bonding between 40 and 44 GPa.

The intensity ratio of the two peaks at the main edge,  $I_1/I_2$ , provides another qualitative probe of changes in the local structure.




**FIGURE 12.** Pressure variation of the intensity ratio ( $I_1/I_2$ ) of the main peaks (see Fig. 9) of  $\text{FeCO}_3$  XANES spectra. The percentage change at the transition pressure is indicated.

The intensity ratio decreases monotonously until 40 GPa, where it then sharply decreases by 18.5% in one step to 44 GPa (Fig. 12). Above 44 GPa, the intensity ratio does not vary significantly with pressure up to 51.5 GPa, the highest pressure reached.

## DISCUSSION

### Spin transition

Changes observed in the Mössbauer, Raman, and XANES spectra of  $\text{FeCO}_3$  that take place between  $\sim 40$  and 45 GPa (or 47 GPa according to Raman data) result from electronic spin crossover of  $3d$  electrons of  $\text{Fe}^{2+}$  atoms. Mössbauer spectra of single-crystal  $\text{FeCO}_3$  show decreasing values of CS with increasing pressure up to 40 GPa that can be explained by increasing  $s$  electron density at the Fe nucleus due to the progressive reduction of interatomic distances. The new component that appears at 40 GPa has zero QS and a CS that is much smaller than the HS component, similar to LS  $\text{Fe}^{2+}$  in ferropentacite and bridgmanite (e.g., Kantor 2007; McCammon et al. 2010). LS  $\text{Fe}^{2+}$  in octahedral coordination has no unpaired electrons with nearly spherical charge distribution; hence the EFG is essentially zero because all  $3d$  energy levels are nearly equally occupied. Spin pairing changes abruptly the degree to which  $s$ -electron density at the Fe nuclei is shielded by  $3d$  electrons; hence the CS drops considerably. The decrease in CS at the transition is roughly 0.3 mm/s, which is higher than the value observed for ferropentacite ( $\sim 0.17$  mm/s, Kantor 2007), but lower than for iron-organic complexes ( $\sim 0.5$  mm/s, Gülich et al. 1978) at ambient or low ( $< 5$  GPa) pressure.

Raman spectroscopy also provides evidence of the spin transition in  $\text{FeCO}_3$ , but through the interatomic vibrations in the sample that are probed by inelastic scattering of light. Unit cell compression between 1 bar and 40 GPa causes a shift of the Raman bands to higher frequencies, to a first approximation following Equation 1

$$\nu = \frac{1}{2\pi} \sqrt{\frac{F_r}{\mu}} \quad (1)$$

where  $\nu$  is the frequency,  $F_r$  is the restoring force, and  $\mu$  is the weighted mass of the molecule or group of atoms.

Between 40 and 47 GPa, the  $A_{1g}$  mode shifts abruptly to lower frequency, comparable to observations by Farfan et al. (2012) and Lin et al. (2012) for natural siderite enriched in manganese and magnesium, respectively. The  $A_{1g}$  mode is attributed to the internal stretching vibration of the carbonate groups. It is therefore strongly dependent on the distance between C and O, and its frequency will decrease with increasing distance. The shift to lower frequency is thus a clear sign of an increase of the C-O distance. This interpretation is consistent with observations by Lavina et al. (2010), who observed an increase in the C-O bond length in natural siderite at 43 GPa and attributed it to structural re-organization caused by spin pairing of Fe<sup>2+</sup> atoms, where shared oxygen atoms that are shared by CO<sub>3</sub>-groups and Fe<sup>2+</sup> are displaced due to the volume collapse of the octahedral site. In contrast, we observed a frequency increase of the external and internal  $E_g$  bands at the spin transition. The external modes, which in FeCO<sub>3</sub> are translational modes between the anion (CO<sub>3</sub>)<sup>2-</sup> and cation (octahedral Fe<sup>2+</sup> coordinated by oxygen) sites, are clearly influenced by the volume collapse of Fe-O octahedra at the transition. The distances between carbonate groups and cations become shorter; hence the frequency increases. The internal  $E_g$  mode, which is caused by in-plane symmetric bending of the carbonate molecule, shifts to higher frequency due to the systematic decrease in the O-O bond length after the transition (Lavina et al. 2010), which induces higher vibrational frequencies due to the decrease of the atomic distance between neighboring oxygen atoms.

Both Mössbauer and Raman data highlight an important point regarding spin pairing in FeCO<sub>3</sub>: under quasi-hydrostatic conditions the process occurs over a pressure interval of at least 5 GPa (Figs. 2 and 6, 7, and 9). The integrated area of the two components in the Mössbauer spectra and the two  $A_{1g}$  bands in Raman spectra allow the percentage of iron atoms in the HS and LS states to be estimated. From Mössbauer spectra at 41 GPa, 12% of Fe<sup>2+</sup> atoms are in the LS state. This observation suggests that the proportion of Fe atoms changing to the LS state does affect the local symmetry and hence the position of the surrounding atoms, inducing short-range (on the angstrom scale) local “structural relaxation” that broadens the transition pressure range. Indeed, since there is a mixture of Fe atoms in HS and LS in the transition region that progressively change to LS, a local structural relaxation that results in stretched bonds between Fe and O will enable some Fe atoms to persist in the HS state over a broader pressure range, being less compressed by the surrounding oxygen atoms.

While a detailed quantitative analysis of XANES spectra incorporating ab initio simulation is beyond the scope of this study, our XANES data can be used to investigate the degree to which spin crossover in FeCO<sub>3</sub> can be detected qualitatively, and how well changes in the local structure can be derived from empirical analysis of the spectra.

The most striking change in the pre-edge region is the abrupt appearance of an additional peak at 7117 eV at 44 GPa. This feature shifted from higher energy, where it was present only as a shoulder in the onset of the main edge. This feature is related to excitations of  $1s$  to  $4p$  electron states of the valence band (e.g., Caliebe et al. 1998), which shift to lower energy with decreasing Fe-O distance. In the actual pre-edge region at ~7112 eV,

which is related to  $1s \rightarrow 3d$  transitions, a change from a doublet feature for HS to a singlet in LS is expected as shown by Westre et al. (1997) by experiment and multiplet analysis on octahedral high- and low-spin complexes. The quality of the data shown here is insufficient for a detailed analysis of the pre-edge due to limited energy resolution and a high level of noise. Nevertheless, a detectable “sharpening” of the pre-edge peak between 40 and 48 GPa can be observed as seen in Figure 9b. Westre et al. (1997) explains the change in the pre-edge region as a change in the accessible electronic states that contribute to the pre-edge after the change in spin state of  $3d$  electrons. In the HS state the octahedral crystal field splits the  $3d$  electron levels into  $e_g$  and  $t_{2g}$  states, giving rise to three possible transitions from the  $1s$  level. Two of the levels are close in energy; hence an apparent doublet is observed for octahedral site symmetry (see also Westre et al. 1997; Wilke et al. 2001). In the LS state only the  $e_g$  level is unoccupied, so only one transition is possible.

In the energy region of the main edge (7120–7140 eV), the peaks in the spectrum of FeCO<sub>3</sub> correspond to a combination of  $1s \rightarrow 4p$  electronic transitions and/or multiple scattering of the photoelectron by neighboring atoms (Fig. 9a, peaks 1 and 2) (Caliebe et al. 1998; Mottana 2004). The positions of these peaks shift to higher energy from 1 bar to 40 GPa (Fig. 10) due to unit-cell compression and the resulting decrease in Fe-O bond length that is also responsible for the increase in the intensity of peak 2 with increasing pressure. The decrease in intensity is likely related to a decrease in the density of  $p$ -like states in that energy region as a consequence of changes in bond length. Between 40 and 44 GPa where spin crossover occurs, the spectra show an abrupt change in the relative amplitudes of peaks 1 and 2, the appearance of peak 3 and the sudden shift to higher energies of the first EXAFS oscillation (roughly from 7170 to 7180 eV, see Figs. 9 and 11). Again, these changes are directly related to the collapse of the Fe coordination polyhedron, i.e., the change in Fe-O distance (Wilke et al. 2007; Bianconi et al. 1983). In addition, the appearance of peak 3 above 44 GPa could arise from multiple scattering of the photoelectron from the neighboring O atoms, which move closer to the Fe atoms after the transition.

While a comprehensive quantitative analysis of the spectra in terms of structural changes would require ab initio calculations that are beyond the scope of this paper, we may use the relation  $\Delta E \times R^2 = \text{const.}$  to estimate the change in the Fe-O distance with pressure from the XANES spectra. To perform this analysis, we use the spectral feature that is located at 7172.5 eV at 0 GPa and define the onset of the edge at the position of the pre-edge (7112 eV) (see also Wilke et al. 2007). This analysis yields  $\Delta E_0 = 60.5$  eV at 0 GPa,  $\Delta E_{40} = 68.9$  eV at 40 GPa, and  $\Delta E_{48} = 75$  eV at 48 GPa. Assuming that the constant in the relation given above is independent of pressure, we can write for 0 and 40 GPa

$$\Delta E_0 \times R_0^2 = \Delta E_{40} \times R_{40}^2, \quad (2)$$

which can be rearranged to

$$(\Delta E_0 / \Delta E_{40})^{0.5} = R_{40} / R_0. \quad (3)$$

From the two values obtained at 0 and 40 GPa, a shortening of the Fe-O distance by a factor of 0.94 is estimated, while across the spin transition from 40 and 48 GPa, we obtain a value of 0.95.

This is comparable to the value of 0.92 obtained by Lavina et al. (2010) based on single-crystal diffraction data (their Fig. 7). The consistency of results is striking, particularly considering the large simplifications made in our analysis, and highlights the sensitivity of XANES to the local environment.

Our results clearly demonstrate the capability of XANES spectra to detect spin crossover in  $\text{FeCO}_3$ . Indeed, the sensitivity has improved markedly since previous studies of other geophysically relevant phases such as ferropericlase (Kantor et al. 2006a) and bridgmanite (Narygina et al. 2009) due to the newly upgraded experimental setup for high-pressure XAFS studies on beamline ID24 at ESRF, together with the implementation of more hydrostatic conditions in the DAC, for example through the use of neon as a quasi-hydrostatic pressure medium. The higher sensitivity could also be related in part to the specific structural arrangement in carbonates [planes of  $(\text{CO}_3)^{2-}$  alternated by octahedrally coordinated cations]. Simulations to quantitatively evaluate the effect of spin crossover on the XAFS spectra are ongoing and will be reported elsewhere.

### Dynamic spin-state distribution

Previous X-ray diffraction (XRD) experiments on siderite and ferromagnesite (e.g., Lavina et al. 2009, 2010; Merlini et al. 2012; Lin et al. 2012) interpreted spin crossover to be a relatively sharp discontinuity taking place at roughly 43 GPa and causing a discontinuous volume reduction of the unit cell. Indeed, XRD is a technique sensitive to long-range structures and cannot probe short-range order such as, for example, HS and LS state distributions. Raman, Mössbauer, and XANES spectroscopy probe the local structure of neighboring atoms and groups of atoms and are therefore sensitive to short-range order features that are not always evident from XRD. Our results, i.e., a successive change of all spectra from HS to LS across a relatively wide pressure range, support the idea of a dynamic nature of spin crossover, which involves a non-static distribution of HS and LS states until the transition is complete. Indeed, a static distribution of the two spin states, which could be considered as a long-range order feature, should be easily resolved by XRD, but so far it has not been reported. Hence, we propose that if a static distribution of the two spin states can be excluded because it was not observed by XRD, the only way to explain the broad and transient pressure range over which spin crossover takes place is a dynamic distribution of HS and LS states in the transition region. On the basis of our result, we suggest that previous XRD studies could be re-examined to investigate if certain features (i.e., the peak splitting and diffuse scattering in Lavina et al. 2010 or the small deviation in the low-pressure equation of state before the volume collapse in Merlini and Hanfland 2013), could reflect a more complex transition.

A direct consequence of the broad spin crossover pressure range is highlighted by the weakening of all Raman  $E_g$  bands as spin crossover is approached, followed by an intensity increase once the final LS state is achieved (Fig. 8). All modes become progressively weaker during compression due to the gradual pressure increase that affects the interaction between the probing laser and the sample, hence reducing signal quality. However, the laser-sample coupling does not explain the almost complete disappearance and reappearance of the lower-frequency modes at the transition. Lavina et al. (2009) described a change in the color of the crystals, where they become progressively darker during and after spin pairing. We observed a similar behavior, where crystals changed from a light reddish color to a slightly darker one. Since we used a green laser with 514 nm wavelength one might attribute the decrease in intensity of the Raman bands to a change in phonon scattering in the visible light region. However, we consider this not to be the case since this does not account for the regain in intensity of the other Raman bands after the transition, and also the internal symmetric stretching vibration  $A_{1g}$  is not affected. Hence, we attribute these intensity minima (Fig. 6) to static disorder caused by spin pairing through the decreased percentage of atoms scattering in the HS state, and consequently the increased amount of scattering by LS atoms. Once the transition is complete, all atoms/molecules in the LS state scatter again equally based on their different vibrational symmetries and new and more intense modes reappear in the spectra. One mode is completely lost after the transition (the external vibration that occurs at  $176 \text{ cm}^{-1}$  at 1 bar); however it is the lowest-frequency  $E_g$  translatory oscillation and might be simply too weak to be detected.

### Grüneisen parameters

We calculated the mode Grüneisen parameters ( $\gamma_i$ ) for each vibrational mode with increase of pressure using the following equation

$$\gamma_i = \frac{K}{\omega} \frac{d\omega}{dP} \quad (4)$$

where the bulk modulus ( $K$ ) is  $117 \pm 1 \text{ GPa}$  (Zhang and Reeder 1999) and  $148 \pm 12 \text{ GPa}$  (Lavina et al. 2010) for the HS and LS states, respectively. We also calculated the pressure shift ( $d\omega/dP$ ) to verify that the differences in the Grüneisen parameters before and after the spin transition were not caused simply by the use of a different bulk modulus (Table 1).

The Grüneisen parameters of HS  $\text{FeCO}_3$  discriminate the two different vibration typologies, internal and external, characterized by smaller and larger values, respectively. In both cases our results match the observations of Santillán and Williams (2004) that were based on infrared measurements. Indeed, the Grüneisen parameters reflect the effect that changing the

**TABLE 1.** Grüneisen parameter ( $\gamma_i$ ) for Raman vibrational modes

| Modes                                                                    | Assignment                      | $d\omega/dP \text{ (cm}^{-1} \text{ GPa}^{-1})$ |      | Mean mode $\gamma_i$ |      |
|--------------------------------------------------------------------------|---------------------------------|-------------------------------------------------|------|----------------------|------|
|                                                                          |                                 | HS                                              | LS   | HS                   | LS   |
| $A_{1g}$ (internal)                                                      | Symmetric (in plane) stretching | 2.17                                            | 1.6  | 0.22                 | 0.21 |
| $E_g$ (internal)                                                         | Symmetric (in plane) bending    | 1.37                                            | 1.86 | 0.21                 | 0.32 |
| $E_g$ (external)                                                         |                                 | 3.82                                            | 2.68 | 1.16                 | 0.72 |
| $E_g$ (external)                                                         |                                 | 2.51                                            |      | 1.18                 |      |
| Mean mode Grüneisen parameter for observed vibrational mode in HS state: |                                 |                                                 |      |                      | 0.69 |
| Mean mode Grüneisen parameter for observed vibrational mode in LS state: |                                 |                                                 |      |                      | 0.42 |

Notes: Bulk moduli from Zhang and Reeder (1998) and Lavina et al. (2009) for HS and LS configuration, respectively. Band assignments from Rutt and Nicola (1974).

volume of a crystal lattice has on its vibrational properties, in this case due to the increase of pressure. The higher values of the external translatory oscillations are consistent with values reported by Ross and Reeder (1992), who described by means of single-crystal XRD the higher compressibility of Fe-O octahedra with respect to the relative incompressibility of the  $\text{CO}_3$  group. Hence, the vibrations associated with the relative translation of the cation with respect to the anionic group are expected to have higher Grüneisen parameters, where the cation site is more affected by the pressure increase due to its higher compressibility.

The Grüneisen parameters measured at spin crossover are also characteristic of the vibrational typology: the internal modes show no variation ( $A_{1g}$ ) or a moderate increase ( $E_g$ ) in their values; whereas the remaining external mode ( $E_g$ ) drops drastically from 1.16 to 0.72, similar to observations by Lin et al. (2012) for ferromagnesite. Indeed at spin pairing the Fe-O octahedra shrink, pulling the oxygen atoms toward the iron atoms and consequently stretching the C-O bonds in the carbonate groups. The external vibration (translatory oscillation) is affected by the shape and the size of the cation site, which is less distorted (Lavina et al. 2010) and has a smaller volume after the transition due to the more compact environment (lower Grüneisen parameter). On the other hand, the stretching of the C-O bond does not influence the carbonate groups; hence the Grüneisen values for the internal vibrations remain constant for symmetric stretching ( $A_{1g}$ ) or show a slight increase for symmetric bending ( $E_g$ ), where the difference is probably related to the different vibrational geometry.

Compared with the study of Lin et al. (2012), we obtain lower values of calculated Grüneisen parameters for both HS and LS states. This may be attributed to the difference in composition and its effect on the bulk modulus, since the latter is the only adjustable parameter used to calculate  $\gamma_i$  (Eq. 4). Indeed, the frequency variation with pressure ( $d\omega/dP$ ) of each mode displays similar values in both studies.

Our calculations relate to the mean mode Grüneisen parameter for all observed vibrations in HS and LS states. Since we recognize that not all phonon modes are taken into account, for example all Raman bands above  $1300\text{ cm}^{-1}$  are excluded, it is not appropriate to compare it with the thermodynamic Grüneisen parameter (e.g., Santillán and Williams 2004; Lin et al. 2012; Liu et al. 2014). Instead our aim is to provide an indication of the difference in compressibility of the  $\text{FeCO}_3$  structure before and after the transition based only on a limited number of vibrations. From our calculations, the average Grüneisen parameters of HS and LS states are, respectively, 0.69 and 0.42. As stated by Lin et al. (2012), the LS state is expected to have a thermodynamic Grüneisen parameter lower than  $\sim 1.1$ , which is the value for the HS state. Our available data suggests the decrease of the Grüneisen parameter to be  $\sim 30\%$ , but this should be tested by further experiments.

#### Effect of stress on Mössbauer spectra collected in a DAC

Differences between powder and single-crystal Mössbauer spectra of  $\text{FeCO}_3$  provide information on the effects of deviatoric and local stresses caused by non-hydrostatic conditions. Any DAC, due to its intrinsic geometry, causes cylindrically

symmetric non-hydrostatic stress fields during compression that affect all materials with non-zero yield strength (e.g., Meng et al. 1993; Dubrovinsky and Dubrovinskaia 2004; Takemura 2001; Zhao et al. 2010). For any type of pressure-transmitting medium there is a pressure limit at which hydrostatic behavior of the medium is lost and consequently deviatoric stresses develop within the sample. These stresses are macroscopic when they are caused by the action of the transmitting medium on the sample and microscopic when they develop at the grain boundaries of polycrystalline samples, i.e., powders (Weidner et al. 1994). Observables from Mössbauer experiments provide qualitative insight into the effects of deviatoric stress on the nuclear behavior and spin crossover of Fe atoms in siderite powder in DACs gas-loaded with neon.

The observed differences between powder and single-crystal Mössbauer spectra arise from the action of microscopic local stresses present at grain boundaries. Indeed, the powder sample was compressed with the intention of producing a pellet with reduced interstitial porosity ( $20\text{ }\mu\text{m}$  size in diameter), so that Ne was acting more as a lubricant between powder grains than as an effective pressure-transmitting medium. This effect is apparent from the influence of pressure on the FWHM of Mössbauer absorption peaks for powder and single-crystals experiments (Fig. 5). With the exception of the 1 bar powder spectrum, all peaks at higher pressures are at least twice as broad for powder compared to single crystals, where the difference increases at higher pressure. The broadening originates from pressure gradients in the gasket hole, which in turn create pressure gradients within the sample. Due to the relatively large cross section of the  $\gamma$ -ray beam in our laboratory ( $\sim 500\text{ }\mu\text{m}$ ) compared to the size of the gasket hole, at each pressure point the resultant Mössbauer spectrum that is measured represents the combination of several spectra collected at slightly higher and lower pressures compared to the pressure measured by ruby fluorescence due to the pressure gradient. In this way spectra characterized by higher and lower CS are merged together and the components in the final spectrum appear broadened. There is little difference in the measured CS of powder and crystals (Fig. 3a), but the differences measured for QS are significant (Fig. 3b). Indeed, the QS values for powder spectra are systematically (except at 1 bar) at least 15% higher than for the single-crystal measurements. The physical explanation is straightforward: powder experiments involve grain-grain interactions that develop under compression and cause the formation of deviatoric stress acting on the local structure around the Fe atoms, causing the Fe-O octahedra to be more distorted than in “regular” quasi-hydrostatic experiments. The distortion induces variations in the shape and size of the charge distribution around the Fe nuclei and ultimately on the electric field gradient, hence on the quadrupole splitting.

The macroscopic and microscopic stresses that develop inside the gasket hole also influence spin crossover. Important differences between powder and single-crystal Mössbauer spectra are: (1) spin crossover starts at lower pressure for single crystals ( $\sim 40\text{ GPa}$ ) compared to powder ( $\sim 50\text{ GPa}$ ); and (2) the pressure range over which spin crossover occurs is smaller for single crystals ( $\sim 5\text{ GPa}$ ) than for powder ( $>10\text{ GPa}$ ). Moreover, the sudden increase ( $\sim 15\%$ ) of QS in powder spectra at the start of spin crossover is caused by enhanced distortion of Fe octahedra

in the crystal structure. Indeed at the spin pairing transition, the dynamic distribution of HS and LS states, with, respectively, larger and smaller octahedra, unbalances the surrounding atomic distribution, stretching or shortening atomic bonds that favor the formation of local stresses.

### IMPLICATIONS

On the basis of this study, we found that spectroscopic techniques provide insight into the short-range atomic order during  $\text{FeCO}_3$  compression from 1 bar to 58 GPa. In addition to XRD, which measures long-range atomic order in crystals, we observed that the spin transition in synthetic  $\text{FeCO}_3$  is a transient process that takes place over a finite range of pressure under quasi-hydrostatic conditions. Moreover, our results demonstrate that spin crossover is a dynamic process during which the Fe atoms in HS and LS states coexist until the LS configuration is fully reached. A static distribution of the two spin states can be excluded based on previous XRD studies that reported only a sudden collapse of unit-cell volume, but no other ordering features.

The calculated Grüneisen parameters for all observed Raman modes agree with the structural arrangement for  $\text{FeCO}_3$  in HS and LS states reported previously in single-crystal XRD studies. While the  $\text{CO}_3$  groups are relatively incompressible over the entire pressure range investigated, Fe-O octahedra show a drastic change in their stiffness after spin crossover, becoming more compact and hence harder to compress.

Deviatoric stress, both macroscopic and microscopic, causes remarkable changes in the high-pressure behavior of  $\text{FeCO}_3$ . Local distortions of the Fe site cause spin crossover to start at higher pressure and a broadening of the spin-transition pressure range.

Our results contribute to understanding the fate of iron-bearing carbonates during subduction. Our experiments clearly show that deformation, which in the Earth is caused by differential stress in the matrix due to processes such as convection, increases the depth at which spin crossover starts and broadens its pressure range beyond 5 GPa. This conclusion is valid both for our room-temperature experiments as well as for mantle conditions. Combined with the results of Liu et al. (2014) who demonstrated that temperature extends the spin crossover region to higher pressure, spin crossover in magnesiosiderite would start at much greater depths (>150 km) than postulated so far in previous work (e.g., Lavina et al. 2009), and would be prolonged over a depth range of at least 300–350 km instead of being a sharp discontinuity in the shallower part of the lower mantle. However, further experiments are required to quantify more precisely the effect of temperature on natural systems under the influence of deviatoric stress.

### ACKNOWLEDGEMENTS

Elite Network Bavaria (ENB), financial support from German Science Foundation (DFG), and Ministry of Research and Education (BMBF) are acknowledged.

### REFERENCES CITED

- Bianconi, A., Dell'Ariccia, M., Gargano, A., and Natoli, C.R. (1983) Bond length determination using XANES. In A. Bianconi, L. Incoccia, and S. Stille, Eds., *EXAFS and Near Edge Structure*, 27, p. 57–61. Springer Series in Chemistry and Physics, Berlin.
- Biellmann, C., Gillet, P., Guyot, F., Peyronneau, J., and Reynard, B. (1993) Experimental evidence for carbonate stability in the Earth's lower mantle. *Earth and Planetary Science Letters*, 118, 31–41.
- Boulard, E., Menguy, N., Auzende, A.L., Benzerara, K., Bureau, H., Antonangeli, D., Corgne, A., Morard, G., Siebert, J., Perrillat, J.P., Guyot, F., and Fiquet, G. (2012) Experimental investigation of the stability of Fe-rich carbonates in the lower mantle. *Journal of Geophysical Research*, 117, B02208.
- Brenker, F.E., Vollmer, C., Vincze, L., Vekemans, B., Szymanski, A., Janssens, K., Szaloki, I., Nasdala, L., Joswig, W., and Kaminsky, F. (2007) Carbonates from the lower part of the transition zone or even the lower mantle. *Earth and Planetary Science Letters*, 260, 1–9.
- Caliebe, W.A., Kao, C.C., Hastings, J.B., Taguchi, M., Kotani, A., Uozumi, T., and de Groot, F.M.F. (1998) 1s2p resonant inelastic X-ray scattering in  $\alpha\text{-Fe}_2\text{O}_3$ . *Physical Review B*, 58, 13452–13458.
- Dasgupta, R., and Hirschmann, M.M. (2010) The deep carbon cycle and melting in Earth's interior. *Earth and Planetary Science Letters*, 298, 1–13.
- Dubrovinsky, L., and Dubrovinskaya, N. (2004) Angle-dispersive diffraction under non-hydrostatic stress in diamond anvil cells. *Journal of Alloys and Compounds*, 375, 86–92.
- Farfan, G., Wang, S., Ma, H., Caracas, R., and Mao, W.L. (2012) Bonding and structural changes in siderite at high pressure. *American Mineralogist*, 97, 1421–1427.
- Franzolin, E., Schmidt, M.W., and Poli, S. (2011) Ternary Ca-Fe-Mg carbonates: Subsolidus phase relations at 3.5 GPa and a thermodynamic solid solution model including order/disorder. *Contributions to Mineralogy and Petrology*, 161, 213–227.
- French, B.M. (1971) Stability relations of siderite ( $\text{FeCO}_3$ ) in the system Fe-C-O. *American Journal of Science*, 27, 37–78.
- Fukao, Y., Obayashi, M., and Nakakuki, T. (2009) Stagnant slab: A review. *Annual Review of Earth and Planetary Science*, 37, 19–46.
- Goldsmith, J.R., Graf, D.L., Witters, J., and Northrop, D.A. (1962) Dolomite-Magnesian calcite relations at elevated temperature and  $\text{CO}_2$  pressures. *Geochimica et Cosmochimica Acta*, 7, 109–128.
- Gütlich, P., Link, R., and Trautwein, A. (1978) *Mössbauer Spectroscopy and Transition Metal Chemistry*. Springer-Verlag, Berlin.
- Harte, B. (2011) Diamond window into the lower mantle. *Science*, 334, 51–52.
- Isshiki, M., Irfune, T., Hirose, K., Ono, S., Ohishi, Y., Watanuki, T., Nishibori, E., Takata, M., and Sakata, M. (2004) Stability of magnesite and its high-pressure form in the lowermost mantle. *Nature*, 427, 60–63.
- Kantor, I. (2007) High-pressure and high-temperature structural and electronic properties of (Mg,Fe)O and FeO, 150 p. Ph.D thesis, University of Bayreuth, Germany.
- Kantor, I., Dubrovinsky, L., McCammon, C., Kantor, A., Pascarelli, S., Aquilanti, G., Crichton, W., Mattesini, M., Almeida, J., Urusov, V.S., and Ahuja, R. (2006a) Pressure-induced phase transition in  $\text{Mg}_{0.80}\text{Fe}_{0.20}$  ferropericlase. *Physics and Chemistry of Minerals*, 33, 35–44.
- Kantor, I., Yu, L., Dubrovinsky, L.S., and McCammon, C. (2006b) Spin crossover in (Mg,Fe)O: A Mössbauer effect study with an alternative interpretation of X-ray emission spectroscopy data. *Physical Review B*, 73, 100101(R).
- Kantor, I., Yu, L., Prakapenka, V., Kantor, A., Dera, P., Kurnosov, A., Sinogeikin, S., Dubrovinskaya, N., and Dubrovinsky, L. (2012) BX90: A new diamond anvil cell design for X-ray diffraction and optical measurements. *Review of Scientific Instruments*, 83, 125102.
- Keppeler, H., Wiedenbeck, M., and Shcheka, S.S. (2003) Carbon solubility in olivine and the mode of carbon storage in the Earth's mantle. *Nature*, 242, 414–416.
- Kurnosov, A., Kantor, I., Boffa-Ballaran, T., Lindhardt, S., Dubrovinsky, L., Kuznetsov, A., and Zehnder, B.H. (2008) A novel gas-loading system for mechanically closing of various types of diamond anvil cells. *Review of Scientific Instruments*, 79, 045110.
- Lavina, B., Dera, P., Downs, R.T., Prakapenka, V., Rivers, M., Sutton, S., and Nicol, M. (2009) Siderite at lower mantle conditions and the effects of the pressure-induced spin-pairing transition. *Geophysical Research Letters*, 36, L23306.
- Lavina, B., Dera, P., Downs, R.T., Yang, W., Sinogeikin, S., Meng, Y., Shenand, G., and Schiferl, D. (2010) Structure of siderite  $\text{FeCO}_3$  to 56 GPa and hysteresis of its spin-pairing transition. *Physical Review B*, 82, 064110.
- Lin, J.F., Liu, J., Jacobs, C., and Prakapenka, V.B. (2012) Vibrational and elastic properties of ferromagnesite across the electronic spin-pairing transition of iron. *American Mineralogist*, 97, 583–591.
- Liu, J., Lin, J.F., Mao, Z., and Prakapenka, V.B. (2014) Thermal equation of state and spin transition of magnesiosiderite at high pressure and temperature. *American Mineralogist*, 99, 84–93.
- Liu, J., Lin, J.-F., and Prakapenka, V.B. (2015) High-pressure orthorhombic ferromagnesite as a potential deep-mantle carbon carrier. *Scientific Reports*, 5, 7640, <http://dx.doi.org/10.1038/srep07640>.
- Mao, H.K., Xu, J., and Bell, P.M. (1986) Calibration of the ruby pressure gauge to 800 kbar under quasi-hydrostatic conditions. *Journal of Geophysical Research*, 91, 4673–4676.
- Mattila, A., Pykkänen, T., Rueff, J.P., Huotari, S., Vankó, G., Hanfland, M., Lehtinen, M., and Härmäläinen, K. (2007) Pressure induced magnetic transition in siderite  $\text{FeCO}_3$  studied by X-ray emission spectroscopy. *Journal of Physics: Condensed Matter*, 19, 382606.
- McCammon, C., Dubrovinsky, L., Narygina, O., Kantor, I., Wu, X., Glazyrin, K., Sergueev, I., and Chumakov, A.I. (2010) Low-spin  $\text{Fe}^{2+}$  in silicate perovskite and a possible layer at the base of the lower mantle. *Physics of the Earth and*

- Planetary Interiors, 180, 215–221.
- Meng, Y., Weidner, D.J., and Fei, Y. (1993) Deviatoric stress in a quasi-hydrostatic diamond anvil cell: Effect on the volume-based pressure calibration. *Geophysical Research Letters*, 20, 1147–1150.
- Merlini, M., and Hanfland, M. (2013) Single-crystal diffraction at megabar conditions by synchrotron radiation. *High Pressure Research*, 33, 511–522.
- Merlini, M., Crichton, W.A., Hanfland, M., Gemmi, M., Mueller, H., Kuzenko, I., and Dubrovinsky, L. (2012) Structures of dolomite at ultrahigh pressure and their influence on the deep carbon cycle. *Proceedings of the National Academy of Sciences*, 109, 13509–13514.
- Merrill, L., and Bassett, W.A. (1975) The crystal structure of  $\text{CaCO}_3$ (II), a high-pressure metastable phase of calcium carbonate. *Acta Crystallographica*, B31, 343–349.
- Mottana, A. (2004) X-ray absorption spectroscopy in mineralogy: Theory and experiment in the XANES region. *EMU Notes in Mineralogy*, 6, 465–552.
- Narygina, O., Mattesini, M., Kantor, I., Pascarelli, S., Wu, X., Aquilanti, G., McCammon, C., and Dubrovinsky, L. (2009) High-pressure experimental and computational XANES studies of  $(\text{Mg,Fe})(\text{Si,Al})\text{O}_3$  perovskite and  $(\text{Mg,Fe})\text{O}$  ferropericlasite as in the Earth's lower mantle. *Physical Review B*, 79, 174115.
- Ono, S., Kikegawa, T., Ohishi, Y., and Tsuchiya, J. (2005) Post-aragonite phase transformation in  $\text{CaCO}_3$  at 40 GPa. *American Mineralogist*, 90, 667–671.
- Pascarelli, S., Mathon, O., Muñoz, M., Mairs, T., and Susini, J. (2006) Energy-dispersive absorption spectroscopy for hard-X-ray micro-XAS applications. *Journal of Synchrotron Radiation*, 13, 351–358.
- Popkov, Yu. A., Eremenko, V.V., Fomin, V.I., and Mokhir, A.P. (1973) Raman scattering of light in antiferromagnetic siderite. *Soviet Physics–Solid State*, 14, 1985–1989.
- Potapkin, V., Chumakov, A.I., Smirnov, G.V., Celse, J.P., Rüffer, R., McCammon, C., and Dubrovinsky, L. (2012) The  $^{57}\text{Fe}$  Synchrotron Mössbauer Source at the ESRF. *Journal of Synchrotron Radiation*, 19, 559–569.
- Prescher, C., McCammon, C., and Dubrovinsky, L. (2012) MossA: a program for analyzing energy-domain Mössbauer spectra from conventional and synchrotron sources. *Journal of Applied Crystallography*, 45, 329–331.
- Ravel, B., and Newville, M. (2005) ATHENA, ARTEMIS, HEPHAESTUS: Data analysis for X-ray absorption spectroscopy using IFEFFIT. *Journal of Synchrotron Radiation*, 12, 537–541.
- Rosenberg, P.E. (1967) Subsolidus relations in system  $\text{CaCO}_3\text{--MgCO}_3\text{--FeCO}_3$  between 350° and 500°C. *American Mineralogist*, 52, 787–796.
- Ross, N.L., and Reeder, R.J. (1992) High-pressure structural study of dolomite and ankerite. *American Mineralogist*, 77, 412–421.
- Rüffer, R., and Chumakov, A.I. (1996) Nuclear Resonance Beamline at ESRF. *Hyperfine Interactions*, 97/98, 589–604.
- Rutt, H.N., and Nicola, J.H. (1974) Raman spectra of carbonates of calcite structure. *Journal of Physics C: Solid State Physics*, 7, 4522.
- Santillán, J., and Williams, Q. (2004) A high-pressure infrared and X-ray study of  $\text{FeCO}_3$  and  $\text{MnCO}_3$ : Comparison with  $\text{CaMg}(\text{CO}_3)_2$ -dolomite. *Physics of the Earth and Planetary Interiors*, 143–144, 291–304.
- Shi, H., Luo, W., Johansson, B., and Ahuja, R. (2008) First-principles calculations of the electronic structure and pressure-induced magnetic transition in siderite  $\text{FeCO}_3$ . *Physical Review B*, 78, 155119.
- Stagno, V., Tange, Y., Miyajima, N., McCammon, C.A., Irifune, T., and Frost, D.J. (2011) The stability of magnesite in the transition zone and lower mantle as function of oxygen fugacity. *Geophysical Research Letters*, 38, L19309.
- Takemura, K. (2001) Evaluation of the hydrostaticity of a helium-pressure medium with powder X-ray diffraction techniques. *Journal of Applied Physics*, 89, 662, <http://dx.doi.org/10.1063/1.1328410>.
- Walter, M.J., Kohn, S.C., Araujo, D., Bulanova, G.P., Smith, C.B., Gaillou, E., Wang, J., Steele, A., and Shirey, S.B. (2011) Deep mantle cycling of oceanic crust: Evidence from diamonds and their mineral inclusions. *Science*, 334, 54–57.
- Weidner, D.J., Wang, Y., and Vaughan, M.T. (1994) Yield strength at high pressure and temperature. *Geophysical Research Letters*, 21, 753–756.
- Westre, T.E., Kennepohl, P., DeWitt, J.G., Hedman, B., Hodgson, K.O., and Solomon, O.E. (1997) A multiplet analysis of Fe K-edge 1s  $\rightarrow$  3d pre-edge features of iron complexes. *Journal of the American Chemical Society*, 119, 6297–6314.
- Wilke, M., Farges, F., Petit, P.E., Brown, G.E. Jr., and Martin, F. (2001) Oxidation state and coordination of Fe in minerals: An Fe K XANES spectroscopic study. *American Mineralogist*, 86, 714–730.
- Wilke, M., Farges, F., Partzsch, G.M., Schmidt, C., and Behrens, H. (2007) Speciation of iron in silicate glasses and melts by in-situ XANES spectroscopy. *American Mineralogist*, 92, 44–56.
- Zhao, J., Angel, R.J., and Ross, N.L. (2010) Effects of deviatoric stresses in diamonds-anvil cell on single-crystal samples. *Journal of Applied Crystallography*, 43, 743–751.
- Zhang, J., and Reeder, R.J. (1999) Comparative compressibilities of calcite-structure carbonates: deviations from empirical relations. *American Mineralogist*, 84, 861–870.

MANUSCRIPT RECEIVED FEBRUARY 2, 2015

MANUSCRIPT ACCEPTED JUNE 23, 2015

MANUSCRIPT HANDLED BY G. DIEGO GATTA